

PRODUCT CATALOGUE





HYALORAL®

Effect of an oral chondroprotective drug in the preventive treatment of elbow dysplasia in Labradors

Simón Martí Angulo, Núria García López. Fundación Once del Perro Guía (FOPG).


OBJECTIVE

- Prevention: To study the incidence of elbow dysplasia in Labrador dogs treated with a prophylactic oral chondroprotective drug
 vs non-treated dogs.
- 2. Treatment efficacy: Clinical follow-up of the animals diagnosed with joint dysplasia (treated vs non-treated).

MATERIAL AND METHODS

Prophylactic and comparative exploratory study in Phase IV.

- 105 dogs (107 joints) of 3-month old Labrador Retrievers.
- Two groups:
 - Control: Specific diet without supplement
 - Treatment: Specific diet with supplement
- Supplement (Hyaloral®, Pharmadiet S.L.U.): tablets composed of hyaluronic acid (20mg), enzymatic hydrolysed collagen (2.2g), glucosamine (500mg), chondroitin sulphate (200mg) and gamma oryzanol (100mg), doses according to weight.
- o 4 follow-up visits at 3, 6, 12 and 20 months.
- o Safety: Follow-up of undesired effects.
- o Clinical evaluation (according to the FOPG protocol)
 - Physical examinations (anamnesis and vital signs)
 - Physiological joint controls (limp, range of movement and swelling)
 - X-ray controls (12 months)
 - ✓ Serological and blood tests (blood count and biochemistry)
 - \checkmark Joint and dysplasia assessments by the educator and veterinarian



PREVENTION RESULTS

- o Dysplasia (at 12 months):
 - ✓ Control group: 35.8% of dogs
 - ✓ Treatment group: 18.5% of dogs


(significantly statistical differences: $p \le 0.05$)

o Affected joints:

- ✓ Elbow: 72.4% of cases.
- √ Hip: 27.6% of cases.
- ✓ Cases of shoulder dysplasia were not diagnosed.

EFFICACY RESULTS

- Animals with radiological diagnosis of joint dysplasia, in both groups.
 Dogs with dysplasia not suitable as guide dogs.
- O Physiological control:
 - ✓ After 12 months, differences between groups (p<0.05) in the most affected joints (left elbow).
 - Significant differences (p<0.05) at 20 months in all the parameters evaluated: limping, reduced mobility and swelling in both joints.
- Physical examination: at visit 3, 88.89% of dogs with joint disorders in the control group vs 40% in the treated group. At the last visit, the difference was 100%.




- Additional controls (blood and serum tests): differences in the values between both groups were not observed.
- Safety: adverse events were not reported in either study group.

O Dysplasia symptoms:

	Treatment				Tatal			
Symptoms		CONTROL (n=19)		HYALORAL (n=10)		Total (n=29)		
dysplasia		N	%	N	%	N	%	р
4.2	No	9	47.4%	8	80%	17	58.6%	0.13
12 months	Yes	10	52,6%	2	20%	12	41.4%	
20 months	No	0	0%	10	100%	10	34.5%	0.00*
20 months	Yes	19	100%	0	0%	19	65.5%	

- * p<0.01, significant differences between groups.
- o Educator evaluation (up to 12 months):
 - Control group: negative assessments in all visits, increasing in each visit up to 33.33%.
 - Treatment group: negative assessments only in the 2nd visit in 40% of the cases, all of which were completely positive in visit 3.



PREVENTION CONCLUSIONS

- Incidence of elbow dysplasia in Labrador dogs is 30% in the normal population.
- o In this study:
- ✓ Incidence of 35.8% was confirmed in the control group.
- Incidence was significantly reduced in the treatment group, up to 18.5%.

Preventive action in the appearance of joint dysplasia in the treatment group vs the control group (95% probability).

EFFICACY CONCLUSIONS IN DYSPLASIA TREATMENT

- o Radiological diagnosis of joint dysplasia at 12 months.
- o Treatment efficacy:
- Limping in the treatment group showed significant improvement vs the control group.
- At 20 months, all the symptoms and signs of dysplasia in the treatment group showed improvement, versus the control group.

Effective treatment for joint dysplasia symptoms.







A comprehensive formula that delivers essential nutrients to **support the proper development and maintenance of joints, tendons, and ligaments**

Instructions for use

Hyaloral® Gel/solution

Kittens: 1.5 ml per day.Adult cats: 3 ml per day.

Dogs:

Up to 5 kg body weight: 1.5 ml per day.
From 5 to 10 kg body weight: 3 ml per day.

Ferrets and rodents: 0.5 ml per kg of body weight per day.

Hyaloral® Small and Medium breeds

Dogs: 1 tablet per 5 kg body weight per day.

Hyaloral® Large and Giants breeds

Dogs: 1 tablet per 20 kg body weight per day in a single dose. • From 40 kg body weight, 2 tablets in a single dose. It is recommended to incorporate **Hyaloral®** to the daily diet for a minimum of 3 months.

Composition per tablet

Hyaloral® Gel/solution per ml:

Enzymatic hydrolyzed collagen 182 mg; Glucosamine 26.9 mg; Chondroitin sulfate 16.6 mg; Gamma oryzanol 8.30 mg; Hyaluronic acid 1.64 mg.

Hyaloral® Small and Medium Breeds per tablet:

Enzymatic hydrolyzed collagen 600~mg; Glucosamine 86.2~mg; Chondroitin sulfate 54.6~mg; Gamma oryzanol 27.3~mg; Hyaluronic acid 5.45~mg.

Hyaloral® Large and Giants Breeds per tablet:

Enzymatic hydrolyzed collagen 2.20 g; Glucosamine 313 mg; Chondroitin sulfate 200 mg; Gamma oryzanol 100 mg; Hyaluronic acid 20.0 mg.

HYALORAL® PLASMA





Hyaloral Plasma 120 tablets Hyaloral Plasma 360 tablets Hyaloral Gel 50 ml Oral use

Indications of use

A comprehensive formula, with nutrients that **support the proper development and maintenance of joints, tendons, ligaments and to support the normal general immune response of the animal**

Instructions for use

Hyaloral® PLASMA Gel/solution

• Kittens: 1.5 ml per day. • Adult cats: 3 ml per day. Dogs:

- Up to 5 kg body weight: 1.5 ml per day.
- 5-10 kg body weight: 3 ml per day.
- Ferrets and rodents: 0.5 ml per kg of body weight per day. Hyaloral® PLASMA

Dogs: 1/4 tablet/5 kg of body weight per day in a single dose. Over 40 kg of body weight, 2 tablets a day in a single dose. Tablet divisible by 4, allowing to dose according to the weight of the pet.

It is recommended to be mixed with the animal's regular food for 3 months minimum. As it is a complementary feed, it can be used continuously.

Composition per tablet

Hyaloral® PLASMA Gel/solution per ml: **Enzymatically hydrolysed collagen** 182 mg; **Glucosamine** 26.9 mg; **Chondroitin sulphate** 16.6 mg; **Plasma proteins** 9.00 mg; **Gamma oryzanol** 8.30 mg; **Hyaluronic acid** 1.64 mg.

Hyaloral® PLASMA per tablet: **Enzymatically hydrolyzed collagen** 2.20 g; **Glucosamine** 313 mg; **Chondroitin sulphate** 200 mg; **Gamma oryzanol** 100 mg; **Plasma proteins** 100 mg; **Hyaluronic acid** 20.0 mg.







Naturally aids the body's response to inflammation, reducing swelling and discomfort.

Instructions for use

Cats: Half a tablet per day.

Dogs:

- Up to 10 kg of body weight: half a tablet per day.
- 10-20 kg of body weight: 1 tablet per day.
- 20-30 kg of body weight: 1.5 tablets per day.
- From 30 kg of body weight: 2 tablets per day.

If preferred, mix with food.

Composition per tablet

Scutellaria baicalensis dried root 100 mg (Baicalin 80.0 mg); Boswellia serrata extract 76.9 mg (Boswellic acids 50.0 mg); Vitis vinifera dried leaves 50.8 mg (Resveratrol 50.0 mg).





Indications of use

Nutritional support for growing animals, to assist in the development of bones and cartilage.

Instructions for use

Cats: 1 tablet per day.

Dogs:

- 1 tablet per 10 kg of body weight per day.
- From 30 kg of body weight: 4 tablets per day.

Composition per tablet

Enzymatic hydrolyzed collagen 265 mg; Calcium 136 mg; Vitamin C 5.10 mg; Folic acid 900 µg; Vitamin D3 2.50 µg.





VETGASTRII





Indications of use

Complementary nutrition supporting gastric imbalances and promoting healthy digestion.

GEL

Cats:

• <5 kg of body weight: 1 ml/day. • >5 kg of body weight: 2 ml/day.

Dogs:

- Up to 5 kg of body weight: 1 ml/day.
- 5-10 kg of body weight: 2 ml/day.
- 10-15 kg of body weight: 3 ml/day.
- 15-20 kg of body weight: 4 ml/day.
- 20-25 kg of body weight: 5 ml/day.

Ferrets, rodents and birds: 0.2 ml per kg of body weight.

Oral route. Shake before use. Take 1-2 hours before the usual feed. If preferred administer half the dose every 12 hours. Animals should always have access to water.

TABLETS

Dogs:

- 1 tablet/20 kg body weight per day.
- From 40 kg body weight, 2 tablets per day.

Oral route. Take 1-2 hours before the usual food. If preferred administer half the dose every 12 hours. Animals should always have access to water.



Includes a 50ml bottle with dosing syringe for oral administration.

Vetgastril® PLUS GEL/SOLUTION per ml:

Aloe barbadensis hydrated leaves 1.00 mg; Hydrolysed keratin 220 mg; Sodium alginate 20.0 mg; Sodium hyaluronate 9.00 mg; Chondroitin sulphate 9.00 mg; Zinc 5.00 mg.

Vetgastril® PLUS TABLETS per tablet:

Hydrolysed keratin 880 mg; Sodium alginate 80.0 mg; Sodium hyaluronate 36.0 mg; Chondroitin sulphate 36.0 mg; Aloe barbadensis dry leaves 20.0 mg; Zinc 20.0 mg.



Includes a 20ml bottle with dosing syringe for oral administration.

Indications of use

A formula enriched with nutrients that support the intestinal system in addressing specific imbalances in dogs and cats.

Instructions for use

Cats:

• Up to 5 kg of body weight: 2 ml/day.

• >5 kg of body weight: 4 ml/day.

• Up to 5 kg of body weight: 2 ml/day.

• 5-10 kg of body weight: 4 ml/day.

• 10-15 kg of body weight: 6 ml/day.

• 15-20 kg of body weight: 8 ml/day.

• From 20 kg of body weight: 10 ml/day.

Shake before use. Administer 2 times a day, dividing the daily amount in two doses.

Composition per ml

Fructo-oligosaccharides (FOS) 98.0 mg; Brewer's yeast (MOS) 98.0 mg; Attapulgite 19.5 mg; Enterococcus faecium 2.00 108 UFC; Potassium chloride 9.00 mg; Sodium chloride 1.50 mg; **Zinc** 0.35 mg.







Natural formula positively supporting the digestive tract and aiding in the elimination of hairballs

Instructions for use

Cats:

• Up to 1 year old: 0.5ml per day.

• Adult cats: 1 ml per day.

Dogs:

• 0.5 ml per 5 kg of weight per day.

• From 35 kg of weight: 4 ml per day. Ferrets and rodents: 0.2 ml per day. Mix with the animal's regular food.

Animals should always have access to water.

It is recommended to follow up after 6 months of use.

Composition per ml

Malt $572 \, \mathrm{mg}$; Fructo-oligosaccharides $86.0 \, \mathrm{mg}$; Yucca shidigera $18.0 \, \mathrm{mg}$ (Saponins $9.00 \, \mathrm{mg}$).





HEPATOSIL® PLUS



Indications of use

A formula enriched with nutrients that support the proper functioning of liver cells

Instructions for use

Hepatosil® PLUS GEL

Dogs and Cats:

1 ml per 5 kg of body weight per day.

Shake before use. It is recommended to be mixed with the animal's regular food.

Water should be available at all times.

It is recommended to follow up after 6 months of use.

Hepatosil® PLUS Small Breeds

Dogs and Cats:

• Up to 7 kg of body weight: 1 tablet per day.

Hepatosil® PLUS Medium Breeds

Dogs:

- From 7 to 10 kg of body weight: 0.5 tablet per day.
- From 10 to 15 kg of body weight: 0.75 tablet per day.
- From 15 to 20 kg of body weight: 1 tablet per day.

Hepatosil® PLUS Large Breeds

Dogs:

- From 20 to 30 kg of body weight: 1 tablet per day.
- From 30 to 35 kg of body weight: 1.5 tablets per day.
- From 35 to 40 kg of body weight: 2 tablets per day.
- From 40 kg of body weight: 2.5 tablets per day.

It is recommended that it be mixed with the animal's regular food. The animals should always have acces to water. It is recommended to follow up after 6 months of use

Composition

Hepatosil® PLUS GEL per ml:

Brewers' yeats 40~mg (Glutation 1.2~mg); Milk thistle extract 45.5~mg (Silybin 15.0~mg); Taurine 100.0~mg; D,L- Methionine 20.0~mg; Vitamin E 6.00~mg; Selenium $6.50~\mu g$; Zinc 3.5~mg; Vitamin B1 $20.0~\mu g$; Folic acid $50.0~\mu g$.

Hepatosil® PLUS Small Breeds per tablet:

Brewers' yeats 312 mg; Vitamin B1 25.0 mg; Thistle seed 49.5 mg (Silybin provided 15.0 mg); Zinc 12.5 mg; Vitamin E 6.00 mg; Folic acid 50.0 µg; Vitamin B12 20.0 µg; Selenium 6.50 µg.

Hepatosil® PLUS Medium Breeds per tablet:

Brewers' yeats $1120\,\text{mg}$; Vitamin B1 $100\,\text{mg}$; Thistle seed $224\,\text{mg}$ (Silybin provided $68.0\,\text{mg}$); Zinc $50.0\,\text{mg}$; Vitamin E $24.0\,\text{mg}$; Folic Acid $200\,\mu\text{g}$; Vitamin B12 $80.0\,\mu\text{g}$; Selenio $26.0\,\mu\text{g}$.

Hepatosil® PLUS Large Breeds per tablet:

Brewers' yeats $1340\,\mathrm{mg}$; Vitamin B1 $120\,\mathrm{mg}$; Thistle seed $297\,\mathrm{mg}$ (Silybin provided $90.0\,\mathrm{mg}$); Zinc $70.0\,\mathrm{mg}$; Vitamin E $32.0\,\mathrm{mg}$; Folic acid $240\,\mathrm{\mu g}$; Vitamin B12 $96.0\,\mathrm{\mu g}$; Selenium $36.0\,\mathrm{\mu g}$.







Nutritional support that aids in the normal functioning of the renal system in dogs and cats

Instructions for use

Cats:

<5 kg of body weight: 1 mldaily.
>5 kg of body weight: 2 ml daily.

Dogs:

- 1 ml / 5 kg body weight per day.
- From 35 kg body weight: 8 ml per day.

Ferrets and rodents: 0.2 ml per day.

Shake before use.
Mix with the usual feed.
Animals should always have access to water.
It is recommended to follow up after 6 months of use.

Composition per ml

Taurine 50.0 mg; Calcium carbonate 45.0 mg; Calcium acetate 40.0 mg; Vitis vinifera dry leaves 25.4 mg (Resveratrol 25.0 mg); Potassium 19.0 mg; Artichocke extract 14.8 mg (Cynarin 740 µg); Salmon oil 6.43 mg (EPA 630 µg; DHA 514 µg); Vitamin B6 330 µg; Folic acid 36.0 µg; Vitamin B12 0.190 µg.





Box containing 60 tablets, in blister packs of 10

Indications of use

Nutritional support formulated with specific nutrients that aid in the normal functioning of the renal system in dogs and cats.

Instructions for use

Cats: 0.5 tablet per day.

Dogs:

- Up to 10 kg of body weight: 0.5 tablet daily.
- 10-20 kg of body weight: 1 tablet daily.
- 20-30 kg of body weight: 1.5 tablets daily.
- Over 30 kg of body weight: 2 tablets daily.

To be taken with food. If preferred, the tablets can be crushed and mixed in with animal regular food.

Animals should always have access to water.

Composition per tablet

Calcium acetate $240 \, \text{mg}$; Vitis vinifera leaves $102 \, \text{mg}$ (Resveratrol $100 \, \text{mg}$); Artichoke extract $60.0 \, \text{mg}$ (Cynarine $3.00 \, \text{mg}$); Green tea dry leaves $20.0 \, \text{mg}$ (EGCG $10.0 \, \text{mg}$); Vitamin B6 $1.30 \, \text{mg}$; Folic acid $134 \, \mu \text{g}$; Vitamin B12 $0.66 \, \mu \text{g}$.







Formula with nutrients that help in the normal maintenance of the urinary system, in particular the lower urinary tract.

Instructions for use

Dogs & cats:

1 ml / 5 kg of of body weight/day. Oral use. Shake before use.

Composition per ml

Glucosamine 100.0 mg; **D-mannose** 90.0 mg; **Cranberry fruits** 60.0 mg; **Horsetail powder** 35.0 mg; **Tryptophan** 15.0 mg; **Chondroitin sulphate** 15.0 mg; **Hyaluronic acid** 8.00 mg.





Contains 30 tablets, in blister strips of 10.

Indications of use

Nutritional support for dogs and cats in **extended-release bilayer tablet**, formulated with nutrients that promote the proper **functioning of the entire urinary system**, **particularly the lower urinary tract**

Instructions for use

Cats: 0.5 tablet / 5 kg of body weight per day. **Dogs:** 1 tablet / 10 kg of body weight per day.

It is recommended to take the tablets whole or half. Do not crush.

Composition per tablet

Bottom layer (immediate release) / Cranberry dried fruits $80.0~\mathrm{mg}$; D-mannose $160~\mathrm{mg}$; Equisetum arvense powder $70.0~\mathrm{mg}$ Top layer (extended release) / Cranberry dried fruits $40.0~\mathrm{mg}$; HC-15 Complex: Sodium hyaluronate $30.0~\mathrm{mg}$; Condroitin sulphate $30.0~\mathrm{mg}$.







Instant formula enriched with nutrients that support the development of kittens from birth to weaning.

Instructions for use

Guidance for the correct development on milk volume necessary for a correct development of the kitten.

(Baby animal age	Cats aged 1-15 days	Cats aged 15-45 days	Ferrets aged 1-7 days	Ferrets aged 7-15 days	Ferrets aged 15-45 days
	Number of daily intakes	6-10	4-6	5	5	5
	Total volume (ml/day)	20-40 ml	40-100 ml	3 ml	4-8 ml	8-12 ml

*From 45 days old until the end of weaning, the number of daily feedings and the quantity of milk substitute will be adjusted based on the infants needs.

Composition per sachet

Linoleic acid $27.1\,\mathrm{mg}$; Linolenic acid $14.1\,\mathrm{mg}$; Omega 3 (EPA+ DHA) $7.50\,\mathrm{mg}$; Colostrum $7.50\,\mathrm{mg}$; Vitamin K $2.83\,\mathrm{\mug}$; Calcium $59.4\,\mathrm{mg}$; Phosphorus $41.9\,\mathrm{mg}$; Vitamin A $133\,\mathrm{UI}$; Vitamin D3 $19.0\,\mathrm{UI}$; Choline chloride $4.50\,\mathrm{mg}$; Taurine: $3.80\,\mathrm{mg}$; Vitamin E $1.90\,\mathrm{mg}$; Niacinamide $1.20\,\mathrm{mg}$; Vitamin C $750\,\mathrm{\mug}$; Calcium D-pantothenate $410\,\mathrm{\mug}$; Vitamin B2 $300\,\mathrm{\mug}$; Vitamin B1 $150\,\mathrm{\mug}$; Vitamin B6 $110\,\mathrm{\mug}$; Folic acid $24.0\,\mathrm{\mug}$; Biotin $7.50\,\mathrm{\mug}$; Vitamin B1 $2.50\,\mathrm{mg}$; Copper $2.50\,\mathrm{mg}$; Copper $2.50\,\mathrm{mg}$; Copper $2.50\,\mathrm{mg}$; Selenium $2.80\,\mathrm{mg}$; Selenium $2.80\,\mathrm{mg}$; Copper $2.50\,\mathrm{mg}$; Copper 2.5





Indications of use

Instant formula enriched with nutrients that support the development of newborn puppies from birth to weaning.

Instructions for use

Guidance for the correct development on milk volume necessary for a correct development of the puppy.

(Baby animal age	Daily volume/100 g of body weight	The number of daily meals	Milk volume /Intake/100 g of body weight	
	1 st week 18-25 ml		6	3-4 ml	
	2 nd week	18-25 ml	5	4-5 ml	
	3 rd week	22-28 ml	3	8 ml	
(From the 4th week*	24-30 ml	3	9 ml	

*From 4 weeks old until the end of weaning, the number of daily feedings and the quantity of milk substitute will be adjusted based on the infants needs.

Composition per sachet

Linoleic acid $35.1\,\mathrm{mg}$; Linolenic acid $18.3\,\mathrm{mg}$; Omega 3 (EPA+ DHA) $3.75\,\mathrm{mg}$; Colostrum $15.0\,\mu\mathrm{g}$; Vitamin K $3.65\,\mu\mathrm{g}$; Calcium $76.8\,\mathrm{mg}$; Phosphorus $54.1\,\mathrm{mg}$; Vitamin A $188\,\mathrm{UI}$; Vitamin D3 $25.0\,\mathrm{UI}$; Choline chloride $5.90\,\mathrm{mg}$; Vitamin E $1.20\,\mathrm{mg}$; Niacinamide $1.20\,\mathrm{mg}$; Vitamin C $750\,\mu\mathrm{g}$; Calcium pantothenate $410\,\mu\mathrm{g}$; Vitamin B2 $300\,\mu\mathrm{g}$; Vitamin B1 $150\,\mu\mathrm{g}$; Vitamin B6 $110\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Biotin $7.50\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Iron $750\,\mu\mathrm{g}$; Manganese $600\,\mu\mathrm{g}$; Zinc $550\,\mu\mathrm{g}$; Copper $110\,\mathrm{mg}$; Selenium $3.80\,\mu\mathrm{g}$







Instant formula enriched with nutrients that support the development of puppies from birth to weaning.

Instructions for use

Guidance for the correct development on milk volume necessary for a correct development of the puppy.

Baby animal age	Daily volume/100 g of body weight	The number of daily meals	Milk volume /Intake/100 g of body weight	
1st week	18-25 ml	6	3-4 ml	
2 nd week	18-25 ml	5	4-5 ml	
3 rd week	22-28 ml	3	8 ml	
From the 4th week*	24-30 ml	3	9 ml	

*From 4 weeks old until the end of weaning, the number of daily feedings and the quantity off milk substitute will be adjusted based on the infants needs.

Composition per sachet:

Linoleic acid $35.1\,\mathrm{mg}$; Linolenic acid $18.3\,\mathrm{mg}$; Omega 3 (EPA+ DHA) $3.75\,\mathrm{mg}$; Vitamin K $3.65\,\mu\mathrm{g}$; Calcium $76.7\,\mathrm{mg}$; Phosphorus $54.1\,\mathrm{mg}$; Sodium $23.7\,\mathrm{mg}$; Vitamin A $188\,\mathrm{UI}$; Vitamin D3 $25.0\,\mathrm{UI}$; Choline chloride $5.90\,\mathrm{mg}$; Vitamin E $1.20\,\mathrm{mg}$; Niacinamide $1.20\,\mathrm{mg}$; Vitamin B2 $300\,\mu\mathrm{g}$; Vitamin B1 $150\,\mu\mathrm{g}$; Vitamin B6 $110\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Biotine $7.50\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Biotine $7.50\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Biotine $7.50\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Polic acid $24.0\,\mu\mathrm{g}$; Polic acid $24.0\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Vitamin B1 $2.300\,\mu\mathrm{g}$; Folic acid $24.0\,\mu\mathrm{g}$; Biotine $3.80\,\mu\mathrm{g}$; Vitamin B1 $3.80\,\mu\mathrm$



CELL HEALTH AND PROTECTION

PROFILAXCEL®



Indications of use

A formula enriched with essential fatty acids and nutrients that support the proper functioning of various organs, including the mammary glands and prostate gland

Instructions for use

Cats: 1 capsule every 2 days.

Dogs

Under 30kg body weight: 1 capsule every 2 days.

Over 30kg body weight: 1 capsule daily.

If you prefer, mix the contents of a capsule with the animal regular feed

Composition per capsule:

Lineseed oil 300 mg (Oleic acid (Omega 9) 52.5 mg, Linoleic acid (Omega 6) 52.5 mg, Linolenic acid (Omega 3)) 173 mg); Pomegranate seed oil 100 mg (Punicic acid (Omega 5)) 70.0 mg); Seaweed oil 87.5 mg (DHA (Omega 3)) 35.0 mg).



LAGRINET® NEO





Indications of use

Hydrating solution for eye hygiene and care.

Instructions for use

Open the bottle and apply one drop to each eye by gently squeezing the container. As it contains no preservatives, it can be used as often as needed. The multi-dose OSD system allows use for up to 3 months after first opening. Suitable for continuous use.

Composition per 100m

Distilled Aloe vera water $10~\rm g$; Distilled Centella asiatica water $10~\rm g$; Sodium hyaluronate $0.15~\rm g$.







Indications of use

Eye solution with a balanced formula that cleans the eye without causing irritation, providing protection against harmful external pollutants.

Instructions for use

Open the bottle and apply one drop to each eye by gently squeezing the container. As it contains no preservatives, it can be used as often as needed. The multi-dose OSD system allows use for up to 3 months after first opening. Suitable for continuous use.

Composition per 100ml

N-Acetylcarnosine 1.00%; Chondroitin sulfate 0.15%; extract *Vaccinium myrtillus* 0.0003%.





Indications of use

Formula based on nutrients that support the animal's ocular system.

Instructions for use

Cats:

- Up to 5 kg of body weight: 1 ml per day.
- Over 5 kg of body weight: 2 ml per day.

Dogs:

- 1 ml per 5 kg in weight per day.
- From 35 kg in weight: 8 ml per day.

Shake before use. Preferably mix with regular feed.

pharmadiet Veterinary

Composition per ml

Omega 3 from fish oil $26.2\,\text{mg}$ (EPA+DHA: $24.2\,\text{mg}$); Taurine $25.3\,\text{mg}$; Algae meal with astaxanthin $13.9\,\mu\text{g}$; Vitamin C $5.17\,\text{mg}$; Dry marigold flowers with lutein $4.12\,\mu\text{g}$; Dry green tea leaves with EGCG $2.06\,\text{mg}$; Dry marigold flowers with zeaxanthin $2.06\,\text{mg}$; Zinc $616\,\mu\text{g}$; D-panthenol $577\,\mu\text{g}$; Dry Vitis vinifera leaves with resveratrol $525\,\mu\text{g}$; Vitamin E $341\,\mu\text{g}$; Lycopene from tomato pulp $308\,\mu\text{g}$; Vitamin A $209\,\mu\text{g}$; Vitamin B6 $209\,\mu\text{g}$; Vitamin B2 $176\,\mu\text{g}$; Vitamin B1 $154\,\mu\text{g}$; Copper $82.5\,\mu\text{g}$; Selenium $20.9\,\mu\text{g}$.







Non-irritant otic solution for external use designed for regular hygiene and cleaning of the ear and ear canal. Helps maintain the physiological conditions of the ear.

Instructions for use

- 1. Position the ear canal as vertically as possible. 2. Apply the product by gently squeezing the container to fill the ear canal with the required amount of NACOTIL®, according to its capacity.
 3. Massage the base of the ear to ensure even distribution of the product. 4. Clean if necessary:
- Use sterile gauze or cotton to gently clean the inside of the ear
- After cleaning, apply 2-3 additional drops of NACOTIL®.
- Wipe the outside of the cannula with clean gauze.

Use NACOTIL® for the maintenance of the physiological conditions of the ear. Wait 30 minutes before any subsequent application. For individual use.

Composition per 100ml

N-Acetyl-cysteine $1.0~\rm g;$ Boric acid $1.4~\rm g;$ Sodium hyaluronate $200~\rm mg;$ Zinc gluconate $2.0~\rm g;$ Citrus medical limonum essential oil $100~\rm mg.$





Indications of use

Otic cleaning solution for external use, non oil based with ph 5.5. Used to regularly clean outer ear. Alcohol free

Instructions for use

Shake the content before use. Squeeze the container and apply the OTINET SOLUTION into the ear canal, whist stretching the ear lobe. Perform a light massage of the ear to allow the correct distribution of the product. If required, clean the inner ear with cotton wool or sterile gauze.

Follow veterinarian instructions.

Composition per 100ml

citrus medical limonum essential oil water 10 g.



OLEODERM®



Pack of 360 capsules in blisters of 10.



Indications of use

A formula enriched with nutrients to support and maintain healthy skin and a shiny coat for dogs and cats.

Instructions for use

Cats: 1 capsule every 2 days.

Dogs:

- Up to 10 kg of body weight: 1 capsule every 2 days.
- From 10 to 15 kg of body weight: 1 capsule per day.
- From 15 to 25 kg of body weight: 2 capsules per day.
- From 25 to 35 kg of body weight: 3 capsules per day.
- Over 35 kg of body weight: 4 capsules per day.

If you prefer, pour the content of a capsule in the animal regular feed. As it's a food supplement, it can be taken continually.

Composition per capsule:

Omega 3 from fish oil 100 mg (EPA 47.5 mg, DHA 32.5 mg); Linoleic acid from borage oil (Omega 6) 95.0 mg; Gammalinolenic acid from borage oil (Omega 6) 50.0 mg; Zinc 5.00 mg; calcium D-pantothenate 1.90 mg; Vitamin B2 286 μg ; Vitamin B1 226 μg ; Folic acid 115 μg ; Biotin 78.0 μg ; Vitamin B1 20.12 μg .





Contains a 50 ml bottle. Gel/solution for topical use.

Indications of use

Moisturising topical gel designed to **protect small skin lesions**, **particularly on extremities and sensitive areas like pads**, **elbows** and **nose**.

Instructions for use

Apply a little amount of **ALOEVET® GEL** and evenly extend on the affected area of the skin. Apply two or three times a day.

Composition per 100ml

Aloe vera gel $32.9~\rm g;$ Centella asiatica extract $3.29~\rm g;$ Hydrolyzed collagen $1.10~\rm g;$ Hyaluronic acid $21.9~\rm mg;$ Vitamin A $13.8~\rm mg;$ Vitamin E $2.36~\rm mg.$







50 ml bottle with syringe dosage syringe. Oral use.

Indications of use

Nutrient-rich formula designed to **support healthy skin and promote a stronger, shinier coat.**

Instructions for use

Cats (< 1 year old): 0.5 ml per day.

Adults cat: 1 ml per day.

Ferrets: 0.5 ml / kg in weight per day.

Shake before use. Mix it with the animal regular feed.

Composition per ml

Malt 559 mg; Taurine 50.0 mg; Salmon oil 26.4 mg; Niacinamide 6.03 mg; Vitamin B5 545 μ g; Folic acid 499 μ g; Biotin 394 μ g; Zinc 302 μ g; Vitamin B6 220 μ g; Vitamin B2 174 μ g; Vitamin B1 162 μ g; Vitamin B12 0.116 μ g.





Box of 60 tablets in blisters of 10

Indications of use

Nutrient-enriched formula designed to support healthy skin and promote a strong, shiny coat in dogs

Instructions for use

1 tablet /10 kg of body weight per day. As it's a food supplement, it can be taken continually.

Composition per tablet

Enzymatically hydrolyzed collagen 199 mg; Hydrolyzed fish collagen 199 mg; Omega 3 from fish oil 20.9 mg; Niacinamide 10.0 mg; Calcium pantothenate 1.00 mg; Folic acid 1.00 mg; Zinc $714 \mu g$; Vitamin B6 273 μg ; Vitamin B2 266 μg ; Vitamin B1 183 μg ; Biotin 9.95 μg ; Vitamin B12 $0.11 \mu g$.



LYSINVIRAL® PLUS





Indications of use

Nutrient-rich formula that supports the normal function of a cat's immune system.

Instructions for use

Cats:

Up to 3 kg of body weight: 1-2 ml daily. 3 kg of body weight or over: 2-4 ml daily.

Shake before use. Take directly or well mixed into the regular

food. Recommended use for 2-3 weeks.

Composition per m

L-Lysine 205 mg; Fish oil 44.8 mg (EPA 15.0 mg, DHA 10.7 mg); Oat bran 12.4 mg (Beta-glucans 8.70 mg); Taurine 15.0 mg; Lactobacillus acidophilus 5.0×10⁷ CFU; Limosilactobacillus reuteri (inactivated) 30.0 mg; Lacticaseibacillus rhamnosus (inactivated) 30.0 mg.





Indications of use

Formula based on nutrients that help the normal functioning of the immune system.

Instructions for use

Cats: 0.5 tablet daily. Dogs and puppies:

- 1 tablet each 10 kg of body weight daily.
- Over 40 kg of body weight 4 tablets daily. Palatable tablets. Can be chewed or crushed.

Composition per tablet

Fresh royal jelly 105 mg; Oat bran 143 mg (Beta glucans 100 mg); Root of *Scutellaria baicalensis* 50.0 mg (Baicalin 40.0 mg); Barley bran 27.9 mg; Ginseng 17.5 mg (Ginsenosides 14.0 mg); Niacinamide 6.00 mg; Calcium pantothenate 2.00 mg; Vitamin B6 680 \mug ; Vitamin B2 570 \mug ; Vitamin B1 460 \mug ; Folic acid 67.0 \mug ; Biotin 50.0 \mug ; Selenium 20.0 \mug ; Vitamin B12 0.400 \mug .

IMMUNITY PLASMORAL®







Box of 60 tablets in blisters of 10

Indications of use

Nutrient-based formula designed to **support the normal functioning of the immune system**, **particularly in situations of immunosuppression**, **before and after vaccinations and deworming**, **enhancing the overall immune response of the animal**.*

Instructions for use

Cats: 0.5 tablet daily.

Dogs and puppies:

- 1 tablet each 10 kg of body weight daily.
- Over 40 kg of body weight 4 tablets daily. Palatable tablets. Can be chewed or crushed.

pharmadiet

Veterinary

Composition per tablet

Plasma proteins 350 mg (Gamma globulins 25.9 mg); Oat bran 143 mg (Beta-glucans 100 mg); Scutellaria baicalensis dry root 50.0 mg (Baicalin 40.0 mg); Ginseng 17.5 mg (Ginsenosides 14.0 mg); Vitamin B6 $670\,\mu g$; Biotin $50.0\,\mu g$; Selenium $20.0\,\mu g$; Vitamin B12 $0.400\,\mu g$.



CALMATONINE XR



Pack containing 120 tablets in blisters of 10 with dispensing envelopes



Indications of use

With nutrients that help to support your pet during times of stress, or anxiety.

Instructions for use

Cats: 0.5 to 1 tablet daily.

Dogs:

- Up to 10 kg of body weight:
- 0.5 to 1 tablet daily.
- From 10–20 kg of body weight: 1 to 2 tablets daily.
- From 20–30 kg of body weight: 1.5 to 3 tablets daily.
- From 30–40 kg of body weight: 2 to 4 tablets daily.
- Over 40 kg of body weight: 3 to 6 tablets daily.

Ferrets and rodents: 0.25 tablet daily.

For occasional use, it is recommended to take 30 minutes to 1 hour before. It can also be taken daily for 6 weeks, continuously. It is recommended to take the tablets whole or split, not crushed.

Composition per tablet

Top layer (immediate release) / Hydrolysed milk protein (provides alpha- casozepin) 80.0 mg Rice germ (provides GABA) 62.5 mg; Valerian root extract 50.0 mg (Valerenic acid 400 µg); Passiflora incarnata extract 50.0 mg (Flavonoids 2.00 mg); Bottom layer (extended release) Tryptophan 130 mg; Vitamin B6 20.0 mg; Magnesium 11.0 mg; Niacinamide 2.77 mg.







Box containing 60 tablets in blisters of 10.



Box containing 30 tablets in blisters of 10.

Indications of use

Nutritional formula designed to support the cognitive system in both dogs and cats.

Instructions for use

Cats: 1 tablet daily.

Dogs:

- Up to 10 kg of body weight: 1 tablet daily.
- From 10-20 kg of body weight: 2 tablets daily.
- From 20-30 kg of body weight: 3 tablets daily.
- From 30–40 kg of body weight: 4 tablets daily.
- Over 40 kg of body weight: 5 tablets daily.

If it is preferred, it can be crushed and mixed with the animal's regular food.

Composition per tablet

Yeast (*Saccharomyces cerevisae*) products with uridine 80.0 mg; Krill meal 75.0 mg (Astaxanthin 5.25 µg, Omega 3 5.06 mg); Tryptophan 50.0 mg; Apple pomace, dried 14.0 mg (Polyphenols 5.60 mg, Quinones 1.40 mg); Biotin 8.00 mg; Zinc 5.00 mg; Vitamin E 2.19 mg; Selenium 6.00 µg.







Nutritional formula designed to enhance the vital tone and energy of the animal.

Instructions for use

Cats:

- Under one year: 0.5 ml daily.
- Over one year: 1 ml daily.

Dogs

- 0.5 ml per 5 kg of body weight per day.
- Over 35 kg: 4 ml per day.

Ferrets: 0.2 mL per 5 kg of body weight per day. Take directly or well mixed into the regular food.

Composition per ml

Fresh royal jelly 170 mg; Taurine 52.5 mg; Salmon oil 6.51 mg; Yucca schidigera 1.21 mg; Tomato pulp 1.13 mg (Lycopene 113 µg); Rice germ 935 µg (Octacosanols 467 µg); Pantothenic acid 0.58 mg; Biotin 0.55 mg; Folic acid 554 µg; Vitamin E 0.41 mg; Vitamin B6 0.25 mg; Beta-carotene 0.22 mg; Vitamin B2 0.19 mg; Vitamin B1 0.17 mg; Selenium 5.01 µg.





Indications of use

Nutritional formula designed to enhance the vitality and energy of dogs, particularly those of advanced age.

Instructions for use

Dogs:


- 1 tablet/10 kg of body weight per day.
- Over 40 kg of body weight, take 4 tablets per day. As it's a food supplement, it can be taken continually.

Composition per tablet

Fresh royal jelly 100~mg; Fish oil (provides Omega 3 (EPA + DHA)) 25.0~mg; Rice germ (provides octacosanols) 17.0~mg; Yucca schidigera (provides saponins) 15.0~mg; Vitamin E 1.60~mg; Vitamin B5 1.00~mg; Tomato pulp $489~\mu g$ (Lycopene $48.9~\mu g$); Folic acid $900~\mu g;$ Vitamin B6 $280~\mu g;$ Vitamin B2 $270~\mu g;$ Vitamin B1 $190~\mu g;$ Beta-carotene $40.0~\mu g;$ Selenium $8.50~\mu g;$ Biotin $8.00~\mu g.$









North & N.I TSR

Email: dhenson-fisher@opko.com Mobile: +44 7428 566152

Francesca

Central TSR

Email: francescagordon@opko.com Mobile: +44 7428 548957

Lucinda

Southwest TSR

Email: lcarter@opko.com Mobile: +44 7428 566016

Samantha

South & Southeast TSR

Email: sfullagar@opko.com Mobile: +44 7428 548967

Kelly

North London TSR

Email: khurworth@opko.com mobile: +44 7428 548853





FOLLOW US!



pharmadiet.uk





GB Third Country Representative & Imported and distributed by:

Pharmadiet UK Ltd. GB154E0797

Unit 2 Leavesden Lodge 1a Leavesden Road, Prohal, Watford, United Kingdom WD24 5FR